

K. Hara

Vol. 44, 1974

No. 2

THE REVIEW OF PHYSICAL CHEMISTRY OF JAPAN

Founded in 1926

CONTENTS

Masaru Nakahara : Derivation of El'yanov and Hamann's Empirical Formula and Another New Formula for Pressure Dependence of Ionization Constants ...	57
Fujio Tanaka : Pressure Effect on Intersystem Crossing from the First Excited Singlet State of Anthracene and Its Derivatives in Solution.....	65
Masami Okamoto : Studies on the Bulk Polymerization of Aliphatic Aldehydes under High Pressure.....	77
Shozo Kishimoto : On the Activity Sequence of Cold-Worked Metal Catalysts for the Hydrogenation of Ethylene.....	89
Tsutomu Takahashi, Kimihiko Hara and Jiro Osugi : The Self-Condensation of Cyclohexanone and Methylcyclohexanone under Very High Pressure.....	93
Tadashi Makita, Yoshiyuki Tanaka and Akira Nagashima : Evaluation and Correlation of Viscosity of Gaseous Ethane and Ethylene.....	98

THE PHYSICO-CHEMICAL SOCIETY OF JAPAN

THE REVIEW OF PHYSICAL CHEMISTRY OF JAPAN

(Butsuri-Kagaku no Shinpo)

Found in 1926

President: Wasaburo Jono

Members of Council:

Michio Okamoto (Chief)

Jiro Osugi

Ryozo Goto

Renpei Goto

Eiji Suito

Hiroyuki Hatano

Board of Editors:

J. Osugi (Chief) *Kyoto University*

R. Goto (Associate) *Kyoto Sangyo University*

E. Suito (") *Kyoto University*

S. Shida *Yokohama University*

H. Teranishi *Kyoto Technical University*

K. Suzuki *Ritsumeikan University*

K. Kuwata *University of Osaka*

S. Seki *University of Osaka*

R. Fujishiro *Osaka City University*

T. Hayakawa *Prefectural University of Osaka*

Y. Kubokawa *Prefectural University of Osaka*

S. Tsuchihashi *University of Kobe*

T. Makita *University of Kobe*

S. Hasegawa *Okayama University*

K. Shimizu *Doshisha University*

H. Togawa *Doshisha University*

Secretary:

M. Sasaki

K. Hara

M. Nakahara

J. Nakaji

April 30, 1975

Communications to the Editor should be addressed to Board of Editors, The Physico-Chemical Society of Japan, Faculty of Science, Kyoto University, Kyoto, Japan.

Business Correspondences should be addressed to: Secretary, The Physico-Chemical Society of Japan, Faculty of Science, Kyoto University, Kyoto, Japan.

Purchase Order should be addressed to: Maruzen Co., Ltd., Nihonbashi, Chuo-ku, Tokyo, Japan.

Published by

THE PHYSICO-CHEMICAL SOCIETY OF JAPAN


(Nippon Butsuri-Kagaku Kenkyu Kai)

Faculty of Science, Kyoto University, Kyoto, Japan

Printed by KAWAKITA PRINTING CO., LTD., Kyoto, Japan

We serve.....
Many peoples,
Many lands,
Many ways.

**SYNTHETIC RESINS
& PLASTICS
INDUSTRIAL
CHEMICALS
FERTILIZERS
DYESTUFFS
AGRICULTURAL
CHEMICALS
MEDICINES
CONSTRUCTION
AND BUILDING
MATERIALS**



**MITSUMI
TOATSU**

MITSUMI TOATSU CHEMICALS, INC.

2-5, Kasumigaseki 3-chome, Chiyoda-ku, Tokyo, Japan

Cable: "MITSUMITOATSU TOKYO"

Telex: Call No. 0222-3622 Answer Back Code (MTCHEM J)

The Review of Physical Chemistry of Japan

Vol. 44, 1974

CONTENTS

No. 1

- Kaoru Date and Hiroji Iwasaki** : Evaluation of P - V - T Properties Data,
The Most Probable Values of Compressibility Factor of Propane and Propene.....1
- Izumi Ishihara, Kimihiko Hara and Jiro Osugi** : Pressure Effects on the
Complexes of Cobalt (II) Chloride and Cobalt (II) Bromide in Acetone Solution.....11
- Tateo Arimoto and Jiro Osugi** : The Charge-Transfer Interaction and Successive
Thermal (2+2) Cycloaddition of α , β - Unsaturated Ethers with Tetracyanoethylene25
- Takeshi Fujii** : Kinetic Studies on the Rearrangement of N -Chloroacetanilide under
Pressure38

No. 2

- Masaru Nakahara** : Derivation of El'yanov and Hamann's Empirical
Formula and Another New Formula for Pressure Dependence of Ionization Constants.....57
- Fujio Tanaka** : Pressure Effect on Intersystem Crossing from
the First Excited Singlet State of Anthracene and Its Derivatives in Solution.....65
- Masami Okamoto** : Studies on the Bulk Polymerization of Aliphatic
Aldehydes under High Pressure77
- Shozo Kishimoto** : On the Activity Sequence of Cold-Worked Metal Catalysts for
the Hydrogenation of Ethylene.....89
- Tsutomu Takahashi, Kimihiko Hara and Jiro Osugi** : The Self-Condensation of
Cyclohexanone and Methylcyclohexanone under Very High Pressure93
- Tadashi Makita, Yoshiyuki Tanaka and Akira Nagashima** : Evaluation and
Correlation of Viscosity of Gaseous Ethane and Ethylene 98


Published by

THE PHYSICO-CHEMICAL SOCIETY OF JAPAN
Faculty of Science, Kyoto University, Kyoto, Japan

HORIBA'S HIGH-PURITY SYNTHETIC CRYSTALS

for precision optical measurements
of radiation, infrared, ultraviolet or laser beam.

NaI (Tl) SCINTILLATOR



HORIBA, Ltd. has established a steady production line for large size crystals of greater than 12 inches by diameter in addition to conventional size crystals

The photograph at right shows an NaI (Tl) Scintillator for use with scintillation camera. The large diameter NaI (Tl) scintillator enabled scintillation survey of an extensive area to be done in a drastically shorter time. HORIBA, Ltd. is ready to respond to your requirements for any purpose as for as the scintillators are concerned.

INFRARED RAY OPTICAL CRYSTAL



Infrared ray optical crystal is formed from materials specially refined for optical crystals as melted and grown into a huge size ingot. Its uncontested high purity, exact crystal lattice and transparency at the transmission range are universally acknowledged. The crystal is available in forms of prism, window or lense of either rough-polish or rough-cut block, or in a random size.

Materials: LiF, NaCl, KCl, KBr, CsI, AgCl, CaF₂, KRS-5, etc.

ELECTRO-OPTIC CRYSTAL



HORIBA, Ltd. is the first in Japan to have developed electro-optic crystal, known as DKDP single crystal especially for modulation of laser beam. The electro-optic effect of the crystal has been greatly expanded by replacing hydrogen in KDP crystal by its isotope, deuterium (maximum deuterium replacement: 99.7%)

Electro-Optical Crystals

	Refractive Index $\lambda = 5461\text{\AA}$	Half Wave Voltage $\lambda = 5461\text{\AA}$	2 Cell Maximum Size
DKDP (D ₂ O, PO ₃ H ₂)	$n_x = 25.4 \pm 10 \text{ m}^{-1}/\text{V}$	3.0KV	40 x 40 x 10mm
LiDP (Li ₂ O, PO ₃ H ₂)	$n_x = 10.3 \pm 10 \text{ m}^{-1}/\text{V}$	7.5KV	40 x 40 x 10mm
LiDP (Li ₂ O, PO ₃ H ₂)	$n_x = 8.1 \pm 10 \text{ m}^{-1}/\text{V}$	9.6KV	40 x 40 x 10mm

HORIBA, Ltd. HORIBA

Miyano Higashi, Kisshoin
Minami-ku, Kyoto, Japan
Phone (075)-313-8121

KANEKA's

products and technologies shall be at your service

KANEKA has now firmly secured its leading position in Japan's chemical industry, and enjoys a high international reputation not only for its capability of offering unique, high-quality products in quick succession but also for its establishment of technical know-how in various field of technology. KANEKA's unique, high-quality products are as follows... "Kanealon," a self-extinguishing modacrylic fiber and is booming for its suitability and nonflammability for carpets, curtains and wigs. "Kane-Ace B," PVC modifier, is now in brisk world-wide demand. "PVC Dispersion Resin," "Heat Resistant PVC," "Expandable Polystyrene," and so on. A variety of epochal projects are now under way at our 10 laboratories far ahead in the same field both at home and abroad.

- * KANEKA's per capita sales and profit are ranked as one of the highest among all chemical manufacturers in the world, thanks to our very high productivity.
- * KANEKA's products have, for the most part, originated in its own laboratories.
- * 55% of KANEKA's products have been developed on its hands for the first time in Japan.

KANEKA



For further details, please write to:

KANEGAFUCHI CHEMICAL IND. CO., LTD.

Head Office:

(Asahi Shinbun Bldg.) 3, 3-chome, Nakanoshima,
Kita-ku, Osaka, Japan
Phone: Osaka (06) 202-1121

Tokyo Office:

(Ohtemachi Bldg.) 4, 1-chome, Ohtemachi, Chiyoda-ku,
Tokyo, Japan
Phone: Tokyo (03) 216-1011

New York Office:

PANAM Bldg., 37th Floor, 200 Park Avenue,
New York, N.Y. 10017, U.S.A.
Phone: (212) 973-4885

Europe Office:

4 Dusseldorf Berliner Allee 26, West Germany
Phone: 12738

